



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\therefore P = \frac{8}{9}F.$$

Assuming \$100 for the face of the note, the proceeds will be \$88 $\frac{8}{9}$, and the discount or interest \$11 $\frac{1}{9}$.

$$\text{From (2), Time} = \frac{\text{Int.}}{P \times r} = \frac{11\frac{1}{9}}{88\frac{8}{9} \times \frac{9}{40}} = \frac{5}{9} \text{ year} = 200 \text{ days.}$$

Also solved by G. B. M. ZERR, and H. C. WHITAKER.

ALGEBRA.

119. Proposed by HARRY S. VANDIVER, Bala, Montgomery County, Pa.

$$\text{Given } \tan x = x + \frac{x^3}{3} + \frac{2x^5}{3 \times 5} + \frac{17x^7}{3^2 \times 5 \times 7} + \frac{62x^9}{3^2 \times 5 \times 7 \times 9} \dots$$

Find the general term and interval of convergence of this series.

I. Solution by COOPER D. SCHMITT, M. A., Professor of Mathematics, University of Tennessee, Knoxville, Tenn.

Using the notation of the Calculus, let $\tan x$ be represented by $f(x)$ or simply f . Then dy/dx will be $f'(x)$ or more simply f' ; similarly, f'' , f''' , f^{iv} , f^v , etc., will be used. We have then by differentiating :

$$\begin{array}{ll} f(x) = \tan x & f(0) = 0. \\ f' = \sec^2 x & f'(0) = 1. \\ f'' = 2 \sec^2 x \tan x = 2ff'' & f''(0) = 0. \\ f''' = 2ff'' + 2f'^2 & f'''(0) = 2. \\ f^{iv} = 6ff'' + 2f''' & f^{iv}(0) = 0. \\ f^v = 6f''^2 + 8ff''' + 2ff^{iv} & f^v(0) = 16. \\ f^{vi} = 20ff''f''' + 10f'f^{iv} + 2ff^v & f^{vi}(0) = 0. \\ f^{vii} = 20f''^2 + 30f'f^{iv} + 12f'f^v + 2ff^{vi} & f^{vii}(0) = 272. \end{array}$$

We see that the *even* differential coefficients vanish and the *odd* follow a remarkable law : The first term is the middle term in the expansion by the Binomial Theorem and the following terms are the double of the successive coefficients of the Binomial Theorem.

By this law we can write any of the odd derivatives, thus :

$$\begin{array}{ll} f^{ix} = 70f^{iv^2} + 112f''f^v + 56f'f^{vi} + 16f'f^{vii} + 2ff^{viii} & f^{ix}(0) = 7936. \\ f^{xi} = 252f^{v^2} + 420f^{iv}f^{vi} + 240f'''f^{vii} + 90f''f^{viii} + 20f'f^x + 2ff^{ix} & f^{xi}(0) = 353792. \\ f^{xiii} = 924f^{vi^2} + 1584f^vf^{vii} + 990f^{iv}f^{viii} + 440f'''f^{ix} + 132f''f^{xi} + 24f'f^{xii} + 2ff^{xiii} \end{array}$$

whence $f^{xiii}(0) = 22368256$, and thus indefinitely.

That is, in writing f^{xiii} , I write the coefficients of $(a+b)^{12}$, which are 1, 12, 66, 220, 495, 792, 924, etc. I set down 924 and double each of the others and get the result as given above.

Substituting the values obtained in MacLaurin's Formula, we have :

$$\begin{aligned}\tan x &= x + \frac{2x^3}{3!} + \frac{16x^5}{5!} + \frac{272x^7}{7!} + \frac{7936x^9}{9!} + \frac{353792x^{11}}{11!} + \frac{22368256x^{13}}{13!} + \text{etc.} \\ &= x + \frac{x^3}{3} + \frac{2x^5}{3 \times 5} + \frac{17x^7}{3^2 \times 5 \times 7} + \frac{62x^9}{3^2 \times 5 \times 7 \times 9} + \frac{1382x^{11}}{3^2 \times 5^2 \times 7 \times 9 \times 11} \\ &\quad + \frac{21844x^{13}}{3^3 \times 5^2 \times 7 \times 9 \times 11 \times 13}.\end{aligned}$$

II. Solution by H. C. WHITAKER, A. M., Ph. D., Professor of Mathematics, Manual Training School, Philadelphia, Pa., and J. SCHEFFER, A. M., Hagerstown, Md.

The general term of the series is

$$\frac{2^{2n}(2^{2n}-1)}{2n!} Bx^{2n-1},$$

where B (Bernoulli) stands for the coefficient of x^n in the expansion of

$$\frac{x}{e^x - 1} = \frac{x}{\frac{x}{1} + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \text{etc.}}$$

120. Proposed by JOSIAH H. DRUMMOND, LL. D., Portland, Me.

A hollow sphere has within it a solid sphere; a quantity of water equal to $1/m$ of the capacity of the hollow sphere is poured in and just covers the solid sphere. Prove that there are two solid spheres, either of which answers the conditions; also find the maximum value $1/m$, beyond which the question is not possible.

I. Solution by J. A. COLSON, Searsport, Me.

The volume of the inner sphere (radius r) + the volume of the water = the volume of a segment whose radius is R and whose height is $2r$; that is

$$\frac{4}{3}\pi r^3 + \frac{4}{3m}\pi R^3 = \frac{1}{3}\pi(2r)^2(3R-2r),$$

which reduces to $r^3 - Rr^2 + R^3/3m = 0 \dots (1)$.

Hence, by Cardan's Method one value of r is

$$\begin{aligned}r_1 &= \frac{1}{3}R + \sqrt[3]{\left[\frac{1}{2}\left[-\frac{R^3}{27}\left(\frac{9}{m}-2\right) + \frac{R^3}{9}\sqrt{\left(\frac{9}{m^2}-\frac{4}{m}\right)}\right]\right]} \\ &\quad + \sqrt[3]{\left[\frac{1}{2}\left[-\frac{R^3}{27}\left(\frac{9}{m}-2\right) - \frac{R^3}{9}\sqrt{\left(\frac{9}{m^2}-\frac{4}{m}\right)}\right]\right]} \dots (2).\end{aligned}$$